arXiv:1302.2834v1 [nucl-th] 12 Feb 2013

Comment on a recent attempt to explain the 'GSI anomaly' by initial-state e.m. spin-rotation coupling*

A. Gal[1]

[1]*Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel*

A recently proposed solution [1] of the 'GSI anomaly' by spin precession of the decaying heavy ions in the magnetic field that controls their circular motion at the GSI storage ring is dubious: the uncertainty in the computed electron-capture decay-rate modulation frequency is at least of order 10^7 Hz, by far exceeding the 1 Hz modulation frequency reported in the GSI experiment [2].

PACS numbers: 13.40.Em, 21.10.Ky, 23.40.-s

Two-body electron-capture decay rates of ^{140}Pr and ^{142}Pm hydrogen-like heavy ions coasting at the GSI Experimental Storage Ring (ESR) were reported to be modulated with period ∼7 s [2]. This modulation is known as the 'GSI anomaly' [3]. Preliminary data exist also for ^{122}I [4]. It was concluded in Refs. [5, 6] that spin precession of the decaying ions in the ESR magnetic field B ≈1.2 T could not give rise to such modulation since the precession frequency associated with the electron is of order $|\mu_e|B \approx 3.3\cdot 10^{10}$ Hz and that associated with the nucleus is of order $\mu_p B \approx 2.5\cdot 10^7$ Hz, where $\mu_{e(p)}$ is the electron (proton) magnetic moment, both these frequencies exceeding by many orders of magnitude the reported modulation frequency $P_{\rm GSI} \sim 1$ Hz. We note that electron-capture decay rates of implanted neutral atoms, not exposed to external magnetic fields, do not show evidence for GSI-like modulation [7, 8].



FIG. 1: $P \equiv P_{\rm LPS} - P_{\rm GSI}$ (Hz units), with $P_{\rm LPS}$ defined by Eq. (1), as a function of the relativistic Lorentz factor $\gamma_{e|n}$. Note that a 0.1% variation of $\gamma_{e|n}$ induces $\approx 4\cdot 10^7$ variation of P. Figure adapted from Ref. [1].

In a recent Letter, however, Lambiase *et al.* [1] noted that a full treatment of spin-rotation coupling results in cancelations among the various components of a spin precession frequency $P_{\rm LPS}$ that splits the two approximately $F=\frac{1}{2}$, $m=\pm\frac{1}{2}$ magnetic states of the rotating ion, bringing $P_{\rm LPS}$ down to the level of $P_{\rm GSI} \sim 1$ Hz. Here,

$$P_{\rm LPS} = a(\gamma_{e|n})\mu_e B + b(\gamma_{e|n}, g_{\rm nucl})\mu_p B, \qquad (1)$$

with coefficients a and b depending on $\gamma_{e|n}$, the relativistic Lorentz factor of the electron with respect to the nucleus, and on the nuclear gyromagnetic ratio $g_{\rm nucl}$. Fig. 1 taken from Ref. [1] shows the dependence of $P \equiv P_{\rm LPS} - P_{\rm GSI}$ on $\gamma_{e|n}$ for ^{140}Pr. To sustain a zero value for P, $(\gamma_{e|n} - 1)$ must be known accurately to at least *three* digits in order to trust the reduction of the μ_e term in Eq. (1) from order 10^{10} Hz down to order 10^7 Hz at which it might get canceled with the μ_p term and, moreover, $g_{\rm nucl}$ must be known accurately to at least *seven* digits so as to reduce the order of magnitude of $P_{\rm LPS}$ from 10^7 Hz further down to $P_{\rm GSI} \sim 1$ Hz. In practice, $(\gamma_{e|n} - 1)$ is estimated semiclassically in Ref. [1] to a single digit only, and the nuclear g.s. gyromagnetic ratio $g_{\rm nucl}$ is either experimentally unknown or known to a single digit only: $g_{\rm nucl}(^{122}{\rm I})=0.94\pm0.03$ [9]. Therefore, the solution proposed in Ref. [1] is dubious.

In conclusion, atomic and nuclear magnetic moments are much too large to resolve the 'GSI anomaly'; see Ref. [10] for a possible, though unlikely solution in terms of a Dirac neutrino magnetic moment of order $10^{-11}\mu_B$.

In memoriam Paul Kienle (1931-2013) who pursued the 'GSI anomaly' with boundless imagination [2–5]; see http://www.lnf.infn.it/edu/stagelnf/2009/interviste09/kienle_ing.htm

---

[1] G. Lambiase, G. Papini, G. Scarpetta, Phys. Lett. B 718 (2013) 998 [arXiv:1205.0684(nucl-th)].
[2] Yu.A. Litvinov *et al.*, Phys. Lett. B 664 (2008) 162.
[3] P. Kienle, Found. Phys. 40 (2010) 733.
[4] P. Kienle, Prog. Part. Nucl. Phys. 64 (2010) 439.
[5] M. Faber *et al.*, J. Phys. G 37 (2010) 015102.
[6] I.M. Pavlichenkov, Phys. Rev. C 81 (2010) 051602.
[7] P.A. Vetter *et al.*, Phys. Lett. B 670 (2008) 196.
[8] T. Faestermann *et al.*, Phys. Lett. B 672 (2009) 227.
[9] V.R. Green *et al.*, Phys. Lett. B 173 (1986) 115.
[10] A. Gal, Nucl. Phys. A 842 (2010) 102.